

INDICATION

DEXTENZA, a corticosteroid (or steroid) medicine, is indicated for treating inflammation (swelling) and pain of the eye following eye surgery.

IMPORTANT SAFETY INFORMATION

Do not use DEXTENZA if you have an active eye infection.

Talk to your doctor if you have a history of high pressure in the eye or glaucoma. Use of steroids for long periods may cause high pressure in the eye. This high pressure may lead to glaucoma, which can harm your vision. Your physician should check your eye pressure during therapy with DEXTENZA.

Talk to your doctor if you have a history of infections of the eye. Steroids may cause an eye infection to worsen or last longer or may increase your risk of infection.

Use of steroids after eye surgery may slow down healing or cause blister formation on the eye.

Tell your doctor if you are pregnant or planning to become pregnant or if you are breastfeeding or planning to breastfeed. The use or effects of DEXTENZA in pregnant women or in the infants of breastfeeding women has not yet been evaluated.

The most common side effects of DEXTENZA were:

- Swelling of the front of the eye, including the iris—the colored area surrounding the pupil—and in the iris-associated muscle (anterior chamber inflammation, including iritis and iridocyclitis)
- Increased pressure in the eye (intraocular pressure increased)
- Problems with seeing clearly (visual acuity reduced)
- Cyst-like swelling in the back of the eye (cystoid macular edema)
- Swelling of the clear front surface of the eye (corneal edema)
- Eye pain
- Redness of the eye (conjunctival hyperemia)
- Headache

These are not all the possible side effects of DEXTENZA. It is important that you tell your doctor right away if you think you may be having any side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch, or call [1-800-FDA-1088](tel:1-800-FDA-1088).